

RESEARCH ARTICLE

Epidemiology and Molecular Characterization of *Cryptosporidium* spp. in Humans, Wild Primates, and Domesticated Animals in the Greater Gombe Ecosystem, Tanzania



CrossMark
click for updates

OPEN ACCESS

Citation: Parsons MB, Travis D, Lonsdorf EV, Lipende I, Roellig DMA, Kamenya S, et al. (2015) Epidemiology and Molecular Characterization of *Cryptosporidium* spp. in Humans, Wild Primates, and Domesticated Animals in the Greater Gombe Ecosystem, Tanzania. PLoS Negl Trop Dis 10(2): e0003529. doi:10.1371/journal.pntd.0003529

Editor: Stephen Baker, Oxford University Clinical Research Unit, VIETNAM

Received: October 31, 2014

Accepted: January 9, 2015

Published: February 20, 2015

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](#) public domain dedication.

Data Availability Statement: All relevant data are within the paper.

Funding: Funding for this study comes from the Morris Animal Foundation (MAF D09ZO-041 and MAF D09ZO-634), the Emory University Global Health Institute, the Arcus Foundation, the Leo S. Guthman Foundation and the National Institutes of Health (R01 AI58715). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Michele B. Parsons^{1,2}, Dominic Travis³, Elizabeth V. Lonsdorf⁴, Iddi Lipende⁵, Dawn M. Anthony Roellig^{2,5}, Shadrack Kamenya⁵, Hongwei Zhang⁶, Lihua Xiao², Thomas R. Gillespie^{1*}

1 Program in Population Biology, Ecology, and Evolution and Departments of Environmental Sciences and Environmental Health, Emory University, Atlanta, Georgia, United States of America, **2** Division of Foodborne, Waterborne, and Environmental Diseases, Centers for Disease Control and Prevention, Atlanta, Georgia, United States of America, **3** College of Veterinary Medicine, University of Minnesota, Minneapolis, Minnesota, United States of America, **4** Department of Psychology, Franklin and Marshall College, Lancaster, Pennsylvania, United States of America, **5** The Jane Goodall Institute, Kigoma, Tanzania, **6** Institute of Parasite Disease Prevention and Control, Henan Center for Disease Control and Prevention, Zhengzhou, China

* thomas.gillespie@emory.edu

Abstract

Cryptosporidium is an important zoonotic parasite globally. Few studies have examined the ecology and epidemiology of this pathogen in rural tropical systems characterized by high rates of overlap among humans, domesticated animals, and wildlife. We investigated risk factors for *Cryptosporidium* infection and assessed cross-species transmission potential among people, non-human primates, and domestic animals in the Gombe Ecosystem, Kigoma District, Tanzania. A cross-sectional survey was designed to determine the occurrence and risk factors for *Cryptosporidium* infection in humans, domestic animals and wildlife living in and around Gombe National Park. Diagnostic PCR revealed *Cryptosporidium* infection rates of 4.3% in humans, 16.0% in non-human primates, and 9.6% in livestock. Local streams sampled were negative. DNA sequencing uncovered a complex epidemiology for *Cryptosporidium* in this system, with humans, baboons and a subset of chimpanzees infected with *C. hominis* subtype IfA12G2; another subset of chimpanzees infected with *C. suis*; and all positive goats and sheep infected with *C. xiaoi*. For humans, residence location was associated with increased risk of infection in Mwamgongo village compared to one camp (Kasekela), and there was an increased odds for infection when living in a household with another positive person. Fecal consistency and other gastrointestinal signs did not predict *Cryptosporidium* infection. Despite a high degree of habitat overlap between village people and livestock, our results suggest that there are distinct *Cryptosporidium* transmission dynamics for humans and livestock in this system. The dominance of *C. hominis*

Competing Interests: The authors have declared that no competing interests exist.

subtype IfA12G2 among humans and non-human primates suggest cross-species transmission. Interestingly, a subset of chimpanzees was infected with *C. suis*. We hypothesize that there is cross-species transmission from bush pigs (*Potamochoerus larvatus*) to chimpanzees in Gombe forest, since domesticated pigs are regionally absent. Our findings demonstrate a complex nature of *Cryptosporidium* in sympatric primates, including humans, and stress the need for further studies.

Author Summary

Cryptosporidium is a common zoonotic gastrointestinal parasite. In a cross-sectional survey of humans, non-human primates (chimpanzees and baboons) and livestock in the Greater Gombe Ecosystem, Tanzania, *Cryptosporidium* infection rate was 4.3%, 16.0% and 9.6% respectively. Infection was not associated with clinical disease in people; however, living in a household with an infected person increased one's risk of infection. Phylogenetic analyses identified clusters of *Cryptosporidium* with a mixed host background. Surprisingly, the Mitumba chimpanzee community, which shares a natural boundary with a human community, had a lower occurrence of *C. hominis* compared to the Kasakela chimpanzee community, which resides in the forest interior (less human exposure). However, Kasakela chimpanzees were also infected with *C. suis*, suggesting a transmission cycle linked to sympatric bush pigs. Our findings highlight the complex nature of zoonotic parasite transmission and stress the need for further studies in similar systems.

Introduction

Cryptosporidium is one of the most important parasitic diarrheal agents in humans in the world, is among the top four causes of moderate-to-severe diarrheal disease in young children in developing nations, and is problematic as an opportunistic co-infection with HIV due to increased morbidity and mortality [1,2]. *Cryptosporidium* is well adapted to zoonotic, water-borne, and foodborne transmission, with a life cycle occurring in suitable hosts and transmission by the fecal-oral route [3]. Zoonoses represent the majority of diseases emerging globally with potential to expand to new host systems, [4] yet despite these health threats, few studies have examined the ecology and epidemiology of this pathogen in rural tropical forest systems characterized by high rates of overlap among humans, domesticated animals, and wildlife [5,6].

In Tanzania, agriculture represents over a quarter of the national income and 80 percent of its labor force [7], but natural resources are declining, affected by desertification and soil degradation from recent droughts. This process has resulted in a high rate of loss of forest and woodland habitat [8]. The resulting fragmented landscape increases human-wildlife contact in these areas, elevating the risk for disease transmission. The Greater Gombe Ecosystem (GGE), Tanzania, in particular, is vulnerable to habitat disturbance, and this has both ecological and financial implications since it is home to diverse wildlife, including endangered chimpanzees (*Pan troglodytes schweinfurthii*), that are important contributors to the national economy through tourism [7].

Gombe National Park, established in 1968, is a small 35-km² forest reserve located 16-km north of Kigoma in Western Tanzania (4°40'S 29°38'E). The park is 1500-m above sea-level with hills sloping westward from a rift escarpment to Lake Tanganyika [9]. It is home to a

number of non-human primate species, including baboons (*Papio anubis*), and a well-known wild chimpanzee population studied continuously for over 50 years [10,11]. There are three chimpanzee communities (Kasekela, Mitumba and Kalande); two of which, (Kasekela, and Mitumba) are habituated [12]. The habitat ranges of these two communities overlap slightly permitting opportunity for member contact. Their habitats have differing degrees of human encroachment [13]. Kasekela, the larger community (~ 65 individuals), is situated at the center of the park in less disturbed forest, whereas Mitumba, the smaller Northern community (~ 25 individuals), is in close proximity to Mwamgongo (4°40'S, 29°34'60' E), a village home to ~ 5000 inhabitants and their livestock. Another village borders the park to the South, but not along the Eastern ridge, due to high elevation and historic soil depletion. Human presence in the park is limited to researchers, tourists, park management staff, local field assistants and members of their families. The park border is not fenced and therefore villagers and their untethered animals (goats, sheep and dogs) are able to enter the park [14]). Mitumba chimpanzees are frequently reported raiding agricultural fields to the east in the Northern village, Mwamgongo, especially during the dry season (I. Lipende, personal communication). There is little evidence that the chimpanzee population has emigrated outside its established habitat for over 20 years, and immigration events are rare.

Death from infectious diseases is the leading cause of mortality for Gombe chimpanzees [15,16]. The chimpanzees have experienced SIVcpz-associated mortality and morbidity, with SIVcpz prevalence ranging between 9–18% and a 10–16-fold higher age-corrected death hazard for infected individuals [17]. *Cryptosporidium* is of special concern in this chimpanzee population, as SIVcpz illness may be complicated by *Cryptosporidium* co-infection, and mirror clinical features observed in human HIV/*Cryptosporidium* co-infections [2], that report *Cryptosporidium* infection rates from 8–30% [18,19]. To improve our understanding of this relationship, and highlight potential management options, we investigated risk factors for *Cryptosporidium* infection and assessed cross-species transmission potential among people, non-human primates, and domestic animals in the GGE, Kigoma District, Tanzania.

Materials and Methods

Ethics statement

This project was reviewed and approved by the Emory University Institutional Review Board (approval #: IRB00018856) under the Expedited review process per 45 CFR 46.110(3), Title 45 CFR Subpart D section 46.404, one parent consent, and 21 CFR 56.110 and the Tanzanian National Institute for Medical Research Institute, Dar Es Salaam, Tanzania, which approved oral consent due to low literacy rates. All adult subjects provided informed consent, and a parent or guardian of any child participant provided informed consent on their behalf. Oral informed consent was obtained by trained local field assistants and documented by witnessed notation on IRB-approved enrollment forms. All animal use followed the guidelines of the Weatherall Report and the NIH Guide for the Care and Use of Laboratory Animals on the use of non-human primates in research, and was approved by the Tanzania Wildlife Research Institute and Tanzania Commission for Science and Technology (permit number 2009-279-NA-2009-184), and the Emory University Animal Care and Use Committee (protocol ID 087-2009). Approval was also obtained from Tanzania National Parks (Permit number TNP/HQ/C10/13) to collect samples from wild chimpanzees. The researchers did not have any interactions with the chimpanzees in the park. All domesticated animals were sampled from households in Mwamgongo village. The owners of the domesticated animals provide verbal consent for the collection of fecal specimens for this study, and the verbal consent was documented. We have

included the GPS coordinates for Mwamgongo village at the first mention of the village in the Introduction.

Sample frame

The study period occurred between March 2010 and February 2011. Paired fecal samples from humans and domestic animals were collected during the dry (July 1–August 15) and wet (November 1–December 15) seasons. Human subjects were either residents of Mwamgongo village (estimated population size (n) ~ 5000) or Gombe National Park (n ~ 100). A baseline demographic survey was performed in June 2010 to identify households within Mwamgongo village with at least one domestic animal species: dog (*Canis lupus*) n ~ 8, goat (*Capra hircus*), n ~ 150 or sheep (*Ovis aries*), n ~ 10. Twenty-five village households with domestic animals were randomly selected for study enrollment. Baboons (n ~ 198) were opportunistically sampled in Mitumba and Kasekela during these two collection periods. Chimpanzees (n ~ 90) were sampled in both Mitumba and Kasekela at quarterly intervals during the course of routine observational health monitoring [16].

Specimen collection and transport

Specimen cups were provided to enrolled village and park residents. Livestock specimens were aseptically collected by a village veterinary officer. Chimpanzee and baboon specimens were non-invasively collected from identified individuals as part of observational health monitoring. All fecal specimens were freshly voided and aseptically transferred to a screw cap plastic vial containing a 2.5% potassium dichromate solution (Fisher Scientific, Pittsburgh, PA). For baboon and non-human primate samples, care was taken to avoid the collection of soil, foliage or water contaminants, by transferring the interior and top most portion of stool to a collection cup using a sterile wooden spatula or swab and avoiding the collection of fecal material in contact with the ground. Each vial was labeled with a unique identification number, and date of collection. Wildlife samples were additionally labeled with the name of the observer, location and animal name. Samples were sealed with Parafilm (Pechiney Plastic Packaging, Chicago, IL) and stored at 4°C, and shipped in ice to Atlanta, GA United States.

Stream sampling

Approximately 1-liter of water was collected in 55-oz sterile Whirl-pak bags (Nasco, Fort Atkinson, WI) and filtered for protozoa using a 0.45-μm Millipore MF-Millipore cellulose ester filter mounted on the Millipore (Billerica, MA) filtration system (diameter 47-mm). When possible, filtration was done on one filter, but in extreme cases where turbidity was high, sequential filtration was performed using two filters. Using sterilized forceps, filters were aseptically transferred to 2-ml cryovials containing a 2.5% potassium dichromate solution. Due to the logistics of field sampling, opportunistic water samples were collected from low, middle and high points of 6 continual streams (dry and wet) and two seasonal streams (wet only). GPS coordinates were obtained using a GPSmap 60CSx from Garmin (Garmin International Inc. Olathe, KS) for each collection point to assist in identifying locations for repeat sampling and if necessary, to assign sampled streams to watersheds associated with specific human or chimpanzee groups.

DNA extraction, molecular detection and subtyping

Nucleic acid was extracted from all fecal specimens and water filters using the FastDNA® SPIN Kit for Soil (MP Biomedicals, LLC, Solon, OH) following the methods described [20]. DNA

extracts were subsequently tested using a polymerase chain reaction and restriction fragment length polymorphism (PCR-RFLP) approach where a segment (~833 bp) of the *Cryptosporidium* SSU rRNA gene is amplified by nested PCR and then species and genotype diagnosis is made by restriction digestion of the secondary PCR product with *Ssp*I (New England BioLabs, Beverly, MA), and either *Vsp*I (Promega, Madison, WI) or *Mbo*II (New England BioLabs) [21,22]. Each sample was run in duplicate by PCR-RFLP analyses with appropriate controls. Specimens that were positive for *Cryptosporidium* by the SSU rRNA PCR were confirmed by DNA sequencing of the 18S PCR products (*C. suis*, *C. xiaoi* and *C. hominis*). A subset of specimens positive for *C. hominis* were also subtyped by sequencing the 60-kilodalton glycoprotein (GP60; ~900 bp) in both directions on an ABI 3130 Genetic Analyzer (Foster City, CA) [23]. All sequences obtained were aligned with reference sequences using MEGA 6.0 or ClustalX software (<http://www.clustal.org/>) to identify *Cryptosporidium* species and *C. hominis* genotypes.

Human risk factor survey

A survey was administered to each human subject focusing on demography, gastrointestinal symptoms (presence or absence within the previous 4 weeks), medication usage, and water usage. To minimize response bias, surveys were administered by trained local field assistants in the national language (Swahili). Data were manually recorded on paper forms, entered into spreadsheets in the computer program Microsoft Excel, and subsequently reviewed for accuracy.

Statistical analyses and control for sample bias

Results were tabulated and compared in Microsoft Excel (Redmond, WA). To control for sample bias we calculated infection rate as the proportion of individuals in each group positive for *Cryptosporidium* divided by the total number of individuals in each group examined [24]. If a single individual sample was positive for *Cryptosporidium*, the subject was considered positive for the collection period (the season for most analyses). Statistical analyses were performed in SPSS version 20.0 (SPSS Inc., Chicago IL). Associations between human survey responses and infection status were compared using logistic regression for categorical binary data. Work history (agricultural fields or forest) were combined as a single factor in the final analysis. Odds ratios (OR) with 95% CI were calculated with significance set at 0.05 for all comparisons. Associations between chimpanzee demographic and observational health data and infection status were evaluated using a generalized estimating equation (GEE) method with exchangeable working correlation structure to account for repeat sampling of individuals. The Huber–White sandwich variance estimation technique was used to calculate confidence intervals (CI). In instances where cells contained less than 5 values, Fisher's exact tests were used to calculate *p*-values.

Results

Six hundred and eighty-four fecal specimens were screened for *Cryptosporidium* including 254 human, 99 domestic animal ($n = 76$ goat, $n = 14$ sheep, $n = 9$ dog) and 331 wildlife ($n = 251$ chimpanzee, $n = 80$ baboon) specimens. *Cryptosporidium* spp. were detected by PCR from 40 (5.8%) fecal samples but was not detected in any water samples ($n = 42$). The infection rate of *Cryptosporidium* was highest among 21/131 (16.0%) nonhuman primates tested, compared to 7/73 (9.6%) livestock and 8/185 (4.3%) humans. No significant differences in frequency were observed between chimpanzees and baboons (Table 1, Fisher's exact test $p = 0.457$) or between the two chimpanzee communities (Table 1, Fisher's exact test $p = 0.7655$). Of the 8 cases of *Cryptosporidium* detected in humans, 7 (87.5%) resided in Mwamgongo village and one

Table 1. Infection rate of *Cryptosporidium* species and *C. hominis* subtypes detected by species and location in and around Gombe National Park, Tanzania.

Host	Positive/Total	<i>Cryptosporidium</i> species detected (n)	<i>C. hominis</i> genotypes	Infection Rate (95% CI)
Humans				
Mwamgongo Village	7/95	<i>C. hominis</i> (7)	IfA12G2 (6/7)	0.07 (0.03–0.15)
Kasekela	0/58			0.0 (0.00–0.08)
Mitumba	1/32	<i>C. hominis</i> (1)	IfA12G2 (1/1)	0.03 (0.03–0.15)
Livestock				
Dog	0/8			0 (0.00–0.40)
Goat	5/56	<i>C. xiaoi</i> (5)		0.09 (0.03–0.20)
Sheep	2/9	<i>C. xiaoi</i> (2)		0.22 (0.03–0.60)
Non-human primates				
Baboon	5/47	<i>C. hominis</i> (5)	IfA12G2 (3/5)	0.11 (0.04–0.24)
Kasekela Chimpanzees				
All <i>Cryptosporidium</i>	12/58			0.21 (0.11–0.34)
<i>C. hominis</i>	6/58 ^a	<i>C. hominis</i> (7)	IfA12G2 (2/7)	0.10 (0.04–0.22)
<i>C. suis</i>	7/58 ^a	<i>C. suis</i> (6)		0.12 (0.05–0.23)
Mitumba Chimpanzees	4/26	<i>C. hominis</i> (4)	IfA12G2 (3/4)	0.15 (0.05–0.36)

^aOne individual was positive for both *C. suis* and *C. hominis*

doi:10.1371/journal.pntd.0003529.t001

(12.5%) in Mitumba camp. No human cases were detected in the Kasekela camp. Sheep had the highest occurrence of *Cryptosporidium* (22%) compared to goat (9%) and dogs (0%) but small sample size prevented evaluation of significance.

We identified three species of *Cryptosporidium* (*C. hominis*, *C. suis* and *C. xiaoi*) in this population (Table 1) based on RFLP and sequence analyses of the SSU rRNA gene. *C. hominis* was detected in all human cases and all 7 cases from sheep and goats were *C. xiaoi*. Six of the 12 positive chimpanzees from Kasekela were genotyped as *C. suis* (a *Cryptosporidium* species predominantly associated with pigs but has been found in a few human cases). This species was not detected in the Mitumba chimpanzee community (Fisher's two-tailed exact test; *p*-value = 0.0537). The porcine species was also not found in the specimens from baboons, humans or domestic animals in the village. The remaining Kasekela chimpanzees (n = 6) and the 4 Mitumba chimpanzees had *C. hominis*. All baboons (n = 5 individuals) also had *C. hominis*. GP60 subtyping of a subset (n = 16) of the *C. hominis* positive samples identified a common subtype IfA12G2 in humans and nonhuman primates. The subtype sequence was identical to two sequences in GenBank; a human *C. hominis* IfA12G2 sequence from South Africa (GenBank accession number JN867334) and a sequence recovered from an olive baboon in Kenya (GenBank accession number JF681172).

We used data from the survey to identify potential risk factors for *Cryptosporidium* infection (Table 2). Among 95 respondents (100%), villagers in Mwamgongo were at greater risk for infection when living with a person who was positive for *Cryptosporidium* (OR = 9.722; 95% CI 1.741–54.279; *p* = 0.011). Persons living with *Cryptosporidium*-positive livestock tended to have a greater odds of infection (OR = 4.750; 95% CI 0.944–23.908; *p* = 0.059). Other factors related to behaviors, including location, occupation (either agricultural or forestry), and not boiling water for consumption were not statistically significant. Although presence of clinical signs was not statistically significant, when reviewing survey data for the *Cryptosporidium* positive patients, 4/8 reported having diarrhea; 2 sought treatment at the village clinic (Flagyl and Paracetemol). Four of 8 households (50%) reported at least one additional

Table 2. Risk factors for *Cryptosporidium* infection in people living in or adjacent to Gombe National Park, Tanzania.

Variable	n	OR	95% CI		p
			Lower	Upper	
Age (≤ 7 years)	179	0.834	0.096	7.277	0.870
Sex (female vs. male)	184	0.946	0.378	2.368	0.905
Seasonality: dry vs wet season	185	2.044	0.244	17.107	0.510
Location (Mwamgongo vs Mitumba)	127	0.406	0.048	3.429	0.407
Location (Mitumba vs Kasekela)	90	5.571	0.2204	140.820	0.356
Location (Kasekela vs Mwamgongo)	153	9.915	0.556	176.950	0.045
Mwamgongo vs park resident	185	7.080	0.853	58.740	0.070
Experienced gastrointestinal symptoms	147	1.458	0.256	8.301	0.671
Used commercial or traditional medicine	173	0.554	0.03	10.194	0.473
Consumption of water from open water source	163	0.156	0.0085	2.875	0.162
Work in agricultural fields or forest	171	3.529	0.665	18.730	0.139
Water not boiled before consumption	153	1.990	0.0992	40.028	0.816
Live in household with another positive person (Mwamgongo)	95	9.722	1.741	54.279	0.011
Live in household with positive livestock (Mwamgongo)	95	4.750	0.944	23.908	0.059

doi:10.1371/journal.pntd.0003529.t002

member of the household experiencing gastrointestinal symptoms, including diarrhea and cramping. Interestingly, affected individuals did not report consuming water from an open source (100%), which appeared protective (OR = 0.156; 95% CI 0.085–2.875) but not statistically significant ($p = 0.162$). 4% of study respondents reported boiling their water before use. No infected individuals reported boiling their drinking water. An association between season and *Cryptosporidium* infection was not observed in humans or non-human primates (Tables 2 and 3). Chimpanzee demographic factors such as age and sex were not risk factors for *Cryptosporidium* infection and evidence of diarrhea was not a reliable predictor of *Cryptosporidium* illness. Kasekela chimpanzees tended to have a higher likelihood (OR = 7.062; 95% CI 0.398–125.251, $p = 0.07$) of infection with *C. suis* as compared to the Mitumba community (Table 3).

Table 3. Risk factors for *Cryptosporidium* infection in chimpanzees in Gombe National Park, Tanzania.

Variable	All <i>Cryptosporidium</i>						<i>C. suis</i>						<i>C. hominis</i>					
	95% CI						95% CI						95% CI					
	n ^a	OR	Lower	Upper	p	OR	Lower	Upper	p	OR	Lower	Upper	p	OR	Lower	Upper	p	
Age (≤ 10 years)	84	1.160	0.395	3.411	0.788	1.950	0.413	9.207	0.399	0.737	0.182	2.992	0.670					
Sex (female vs. male)	84	0.425	0.128	1.412	0.162	0.181	0.022	1.504	0.113	0.716	0.172	2.980	0.647					
Seasonality: (dry vs wet)	84	1.194	0.440	3.245	0.728	0.875	0.226	3.391	0.848	1.471	0.459	4.745	0.514					
Location (Kasekela vs Mitumba)	84	0.935	0.274	3.187	0.915	7.062	0.398	125.251	0.070	2.354	0.616	8.998	0.211					
Observed to have diarrhea	79	2.034	0.703	5.883	0.190	1.461	0.264	8.069	0.664	2.425	0.632	9.309	0.197					

^aSample sizes may vary based on number of individual observations.

Binary logistic regression was used to calculate odds ratios, confidence interval and significance in most cases. Fisher's Exact test was used to calculate p-values when cells contained values less than 5.

doi:10.1371/journal.pntd.0003529.t003

Discussion

Of 90 humans residing in camps within Gombe National Park, only one *Cryptosporidium* infection was observed (1%). In contrast, *Cryptosporidium* infection in residents of Mwamgongo village reached 10% during the drier months. These frequencies are comparable to those reported in some studies from children and adults without HIV, ranging from 0–18% [25–27], though frequencies as high as 32% have been reported elsewhere [28]. Site and HIV prevalence would seem to be important factors in human occurrence rates. Although HIV testing was beyond the scope of this study, a recent country report [9] indicates a low HIV prevalence (< 1%) from the Kigoma region where the study occurred. Infection was not statistically associated with gastrointestinal illness or stool consistency for humans or wildlife, findings consistent with earlier studies [28–32]. Surprisingly, unsafe drinking water (i.e. untreated/unboiled water, open water source) was not found to increase risk of *Cryptosporidium* infection, which could be the result of degradation or improper tapping of the water line, exposing water to environmental contamination [33]. This warrants further study as consumption of contaminated ground water has been repeatedly associated with *Cryptosporidium* infection [34,35].

Cryptosporidium was detected in baboons (11%) and the two chimpanzee communities (15–21%) at moderate frequencies. Similar frequencies of *Cryptosporidium* have been detected previously in other African primates. Habituated mountain gorillas in Bwindi Impenetrable National Park (BINP), Uganda had *Cryptosporidium* in (11%) of specimens sampled; 73% of positive specimens were detected from human-habituated gorillas [36]. Genetic characterization of this population determined that the gorillas and the local human community both carry *C. parvum* [37]. Eleven percent of red colobus and black-and-white colobus in Kibale National Park (KNP), Uganda were infected with *Cryptosporidium*; genetic sequences from some humans and colobus from KNP were identical, while two strains from red colobus in the forest interior were infected with a divergent subclade, suggesting the possibility of separate zoonotic and sylvatic cycles [28]. These findings support the zoonotic spillover potential of *Cryptosporidium* among humans, wildlife and livestock.

Similar patterns have been observed with other directly or environmentally transmitted enteric pathogens. At KNP, *Giardia* was detected in red colobus in forest fragments (5.7%), but not detected in undisturbed forest (0%) [38]. Genetic analyses of recovered strains determined that red colobus were infected with *Giardia duodenalis* assemblages associated with humans and livestock, suggesting complex cross-species transmission [39]. *Escherichia coli* strains from gorillas in BINP and chimpanzees in KNP that overlapped with humans were more similar to strains collected from resident humans and livestock compared to strains collected from gorillas and chimpanzees living in undisturbed forest [40,41]. Additionally, in KNP, genetic similarity between human/livestock and primate bacteria increased three-fold with a moderate to high increase in anthropogenic disturbance of forest fragment. Bacteria harbored by humans and livestock were more similar to those of monkeys that entered human settlements to raid crops, than to bacteria of other primate species [42]. These findings reinforce the notion that habitat overlap and anthropogenic disturbance increase the risk of interspecies transmission between wildlife, humans and livestock and that transmission can occur both via direct physical contact with or ingestion of contaminated feces and by indirect exposure via a shared (potentially contaminated) watershed. Although water samples screened in this study were negative for the parasite, waterborne outbreaks of *Cryptosporidium* as a result of human and animal fecal contamination are common [43–45]. Watershed sampling for *Cryptosporidium* in this study was opportunistic using smaller volumes of water than advocated by standard screening protocols [46] and provided only limited inter-seasonal sampling [47–49]. Therefore, our negative results do not assure that waterborne transmission is not important in this system. Future studies

using more comprehensive watershed sampling would help to resolve this aspect of *Cryptosporidium* transmission.

The results of our RFLP and sequence analyses of the SSU rRNA gene suggest multiple potential zoonotic pathways for *Cryptosporidium* transmission in this study system. The village data reinforces our understanding that species of *Cryptosporidium* vary in their zoonotic potential [50]. Our data suggest that there is less likelihood that the *C. xiaoi* affecting the livestock is capable of causing illness in humans, considering the close proximity of livestock to humans in this community [i.e., animals often residing in homes], where animal-human contact is quite high though it has been documented in other studies [51]. *Cryptosporidium hominis* was also not detected among the domesticated animals in this study, but *C. hominis* has been found in other studies to be a zoonotic species, affecting both humans and domesticated animals [52,53]. Homes with positive livestock had a tendency for increased risk of human infection suggesting contribution of environmental factors or behaviors that may place the household at increased risk.

We anticipated that the Kasekela chimpanzee community would have a lower occurrence of *Cryptosporidium* compared to Mitumba, which shares a natural border with Mwamgongo village. Although not statistically significant, there was a higher frequency of *Cryptosporidium* recovered in Kasekela. The occurrence of *C. hominis* was not statistically higher for Mitumba compared to Kasekela. However, the results demonstrated that 50% (6/12) of the *Cryptosporidium* recovered from the Kasekela community are *C. suis*, a species associated with pigs. Domestic pigs are not found in the park or village due to religious preference (predominantly Muslim communities). However, bush pigs (*Potamochoerus larvatus*), native to the Gombe forest habitat, are common in Kasekela but have not been observed frequently in Mitumba forest. Thus bush pigs may serve as the reservoir host for *C. suis* in this system. Chimpanzees may be infected through contact with this animal (i.e., chimpanzees occasionally consume bush pigs) or by indirect contact with infective feces on the forest floor or the contamination of shared water sources. This putative pathway of transmission is supported by the fact that *C. suis* was not recovered in the Mitumba chimpanzee community, the baboons, domesticated livestock or village inhabitants. *C. suis* has zoonotic potential having been previously identified in an HIV+ patient in Lima, Peru, [54] and from patients in Henan, China and England [55,56]. Similarly, Salyer et al [28] found that while *Cryptosporidium* may be transmitted frequently among domesticated animals, humans and wildlife in areas of overlap, there may be host-parasite specific dynamics that occur in the absence of these regular interactions creating separate transmission cycles.

The appearance of the *C. hominis* species and its subtype IfA12G2C among the humans, baboons and chimpanzee communities demonstrates the zoonotic transmission potential of this parasite species among these closely related host species and points to a dominance of anthropozonotic transmission in this system. This subtype has been previously reported among captive olive baboons in Kenya [57], and is prevalent in humans in Africa [58]. Additionally, common primate behaviors may increase the likelihood for animal to human transmission. For example, baboons raid camp food reserves and homes, potentially transmitting etiologic agents via infected feces to humans. The Mitumba chimpanzees are reported to raid agricultural fields just outside the park boundaries, which can transmit diseases from the potentially contaminated feces of livestock or exposed human sewage.

Our results are based on small sample sizes, that if increased could alter the frequency and predictions of *Cryptosporidium* subtypes, infection and illness. The finding of *C. suis* in the Kasekela chimpanzee community is novel. We presume these chimpanzees experienced cross-species transmission from bush pigs in Gombe forest, because domesticated pigs are absent from the area. Unfortunately, we do not have access to fecal specimens from the local bush pig

population to compare to specimens recovered from Kasekela chimpanzees. Despite the high overlap observed between people and livestock in villages in this region, our results suggest that the transmission dynamics of *Cryptosporidium* for humans and livestock are distinct but the dominance of *C. hominis* in humans and non-human primates suggest the potential for cross-species transmission. Our findings highlight the complex nature of zoonotic parasite transmission and stress the need for further studies in similar systems.

Supporting Information

S1 Checklist. STROBE Checklist.
(PDF)

Acknowledgments

We are grateful to J. Baranyikwa and the field assistants of Gombe Stream Research Centre for assisting in the collection of chimpanzee health data and fecal specimens. We thank the Mwangongo Village Animal Health Officer, E. Mchalo for assisting in the collection of domestic animal specimens. We also thank the Kigoma district health officers and D. Mjungu for discussion and the Tanzanian Commission for Science and Technology, Tanzania National Parks, and Tanzania Wildlife Research Institute for permission to conduct the research.

The authors would like to warmly acknowledge T. Dearen for technical expertise, A. Blackstock for statistical guidance, and J. Bodager and K. Cross for laboratory assistance.

Author Contributions

Conceived and designed the experiments: TRG MBP. Performed the experiments: TRG MBP DT EVL IL LX DMR HZ SK AC. Analyzed the data: TRG MBP EVL DMR LX. Wrote the paper: TRG MBP.

References

1. Kotloff KL, Nataro JP, Blackwelder WC, Nasrin D, Farag TH, et al. (2013) Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *Lancet* 382: 209–222. doi: [10.1016/S0140-6736\(13\)60844-2](https://doi.org/10.1016/S0140-6736(13)60844-2) PMID: [23680352](https://pubmed.ncbi.nlm.nih.gov/23680352/)
2. Tzipori S (1988) CRYPTOSPORIDIOSIS IN PERSPECTIVE. *Advances in Parasitology* 27: 63–129. PMID: [3289331](https://pubmed.ncbi.nlm.nih.gov/3289331/)
3. Smith HV, Caccio SM, Cook N, Nichols RAB, Tait A (2007) *Cryptosporidium* and *Giardia* as foodborne zoonoses. *Veterinary Parasitology* 149: 29–40. PMID: [17728067](https://pubmed.ncbi.nlm.nih.gov/17728067/)
4. Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, et al. (2008) Global trends in emerging infectious diseases. *Nature* 451: 990–993. doi: [10.1038/nature06536](https://doi.org/10.1038/nature06536) PMID: [18288193](https://pubmed.ncbi.nlm.nih.gov/18288193/)
5. Leav BA, Mackay M, Ward HD (2003) *Cryptosporidium* species: New insights and old challenges. *Clinical Infectious Diseases* 36: 903–908. PMID: [12652391](https://pubmed.ncbi.nlm.nih.gov/12652391/)
6. Appelbee AJ, Thompson RCA, Olson ME (2005) *Giardia* and *Cryptosporidium* in mammalian wildlife—current status and future needs. *Trends in Parasitology* 21: 370–376. PMID: [15982929](https://pubmed.ncbi.nlm.nih.gov/15982929/)
7. US Counter Intelligence Agency (2013) "Tanzania". *The World Factbook*.
8. Ministry of Agriculture Food Security and Cooperatives. The United Republic of Tanzania (2008) <http://kilimo.go.tz/agricultural%20maps/Tanzania%20Soil%20Maps/Soil%20maps.htm>.
9. Wallis J, Lee DR (1999) Primate conservation: The prevention of disease transmission. *International Journal of Primatology* 20: 803–826.
10. Goodall J (1986) *The Chimpanzees of Gombe: Patterns of Behavior*. Boston: Belknap Press of the Harvard University Press PMID: [25101439](https://pubmed.ncbi.nlm.nih.gov/25101439/)
11. Earnhardt JM, Rossi SR, Lonsdorf EV, Pusey AE (2003) A demographic comparison of wild chimpanzees from Gombe and a managed population from North American zoos. *American Journal of Primatology Suppl.* 60:62–63.

12. Pusey AE, Pintea L, Wilson ML, Kamenya S, Goodall J (2007) The contribution of long-term research at Gombe National Park to chimpanzee conservation. *Conservation Biology* 21: 623–634. PMID: [17531041](#)
13. Pusey AE, Wilson ML, Collins DA (2008) Human impacts, disease risk, and population dynamics in the chimpanzees of Gombe National Park, Tanzania. *American Journal of Primatology* 70: 738–744. doi: [10.1002/ajp.20567](#) PMID: [18521891](#)
14. Parsons MB, Gillespie TR, Travis D, Lipende I, Gilagiza B, Kamenya S, Pintea L, Vazquez-Prokopen GM (2014) Global Positioning System Data-Loggers: A Tool to Quantify Fine-Scale Movement of Domestic Animals to Evaluate Potential for Zoonotic Transmission to an Endangered Wildlife Population. *PLoS ONE* 9(11): e110984. doi: [10.1371/journal.pone.0110984](#) PMID: [25365070](#)
15. Williams JM, Lonsdorf EV, Wilson ML, Schumacher-Stankey J, Goodall J, et al. (2008) Causes of death in the Kasekela chimpanzees of Gombe National Park, Tanzania. *American Journal of Primatology* 70: 766–777. doi: [10.1002/ajp.20573](#) PMID: [18506732](#)
16. Lonsdorf EV, Travis D, Pusey AE, Goodall J (2006) Using retrospective health data from the Gombe chimpanzee study to inform future monitoring efforts. *American Journal of Primatology* 68: 897–908. PMID: [16900499](#)
17. Keele BF, Jones JH, Terio KA, Estes JD, Rudicell RS, et al. (2009) Increased mortality and AIDS-like immunopathology in wild chimpanzees infected with SIVcpz. *Nature* 460: 515–519. doi: [10.1038/nature08200](#) PMID: [19626114](#)
18. Tarimo DS, Killewo JZ, Minjas JN, Msamanga GI (1996) Prevalence of intestinal parasites in adult patients with enteropathic AIDS in north-eastern Tanzania. *East African medical journal* 73: 397–399. PMID: [8840602](#)
19. Colebunders R, Lusakumuni K, Nelson AM, Gigase P, Lebughe I, et al. (1988) Persistent diarrhoea in Zairian AIDS patients: an endoscopic and histological study. *Gut* 29: 1687–1691. PMID: [3220308](#)
20. da Silva AJ, Bornay-Llinares FJ, Moura INS, Slemenda SB, Tuttle JL, et al. (1999) Fast and reliable extraction of protozoan parasite DNA from fecal specimens. *Molecular Diagnosis* 4: 57–64. PMID: [10229775](#)
21. Xiao LH, Bern C, Limor J, Sulaiman I, Roberts J, et al. (2001) Identification of 5 types of *Cryptosporidium* parasites in children in Lima, Peru. *Journal of Infectious Diseases* 183: 492–497. PMID: [11133382](#)
22. Xiao L, Ryan U (2008) *Cryptosporidium* and *Cryptosporidiosis*; Fayer R, Xiao L, editor. New York, USA: CRC Press. PMID: [25506952](#)
23. Alves M, Xiao LH, Sulaiman I, Lal AA, Matos O, et al. (2003) Subgenotype analysis of *Cryptosporidium* isolates from humans, cattle, and zoo ruminants in Portugal. *Journal of Clinical Microbiology* 41: 2744–2747. PMID: [12791920](#)
24. Gillespie TR, Lonsdorf EV, Canfield EP, Meyer DJ, Nadler Y, et al. (2010) Demographic and Ecological Effects on Patterns of Parasitism in Eastern Chimpanzees (*Pan troglodytes schweinfurthii*) in Gombe National Park, Tanzania. *American Journal of Physical Anthropology* 143: 534–544. doi: [10.1002/ajpa.21348](#) PMID: [20623606](#)
25. Cegielski JP, Ortega YR, McKee S, Madden JF, Gaido L, et al. (1999) *Cryptosporidium*, Enterocytozoon, and *Cyclospora* Infections in Pediatric and Adult Patients with Diarrhea in Tanzania. *Clinical Infectious Diseases* 28: 314–321. PMID: [10064250](#)
26. Tumwine JK, Kekitiinwa A, Bakeera-Kitaka S, Ndeezi G, Downing R, et al. (2005) Cryptosporidiosis and microsporidiosis in Ugandan children with persistent diarrhea with and without concurrent infection with the human immunodeficiency virus. *American Journal of Tropical Medicine and Hygiene* 73: 921–925. PMID: [16282304](#)
27. Gomez Morales MA, Atzori C, Ludovisi A, Rossi P, Scaglia M, et al. (1995) Opportunistic and non-opportunistic parasites in HIV-positive and negative patients with diarrhoea in Tanzania. *Tropical medicine and parasitology: official organ of Deutsche Tropenmedizinische Gesellschaft and of Deutsche Gesellschaft fur Technische Zusammenarbeit (GTZ)* 46: 109–114.
28. Salyer SJ, Gillespie TR, Rwego IB, Chapman CA, Goldberg TL (2012) Epidemiology and Molecular Relationships of *Cryptosporidium* spp. in People, Primates, and Livestock from Western Uganda. *Plos Neglected Tropical Diseases* 6.(4):e1597. doi: [10.1371/journal.pntd.0001597](#) PMID: [22506085](#)
29. Cama VA, Ross JM, Crawford S, Kawai V, Chavez-Valdez R, et al. (2007) Differences in clinical manifestations among *Cryptosporidium* species and subtypes in HIV-infected persons. *J Infect Dis* 196: 684–691. PMID: [17674309](#)
30. Checkley W, Gilman RH, Epstein LD, Suarez M, Diaz JF, et al. (1997) Asymptomatic and Symptomatic Cryptosporidiosis: Their Acute Effect on Weight Gain in Peruvian Children. *American Journal of Epidemiology* 145: 156–163. PMID: [9006312](#)

31. Houpt ER, Bushen OY, Sam NE, Kohli A, Asgharpour A, et al. (2005) Short report: Asymptomatic *Cryptosporidium hominis* infection among human immunodeficiency virus-infected patients in Tanzania. American Journal of Tropical Medicine and Hygiene 73: 520–522. PMID: [16172475](#)
32. Gracenea M, Gómez MS, Torres J, Carné E, Fernández-Morán J (2002) Transmission dynamics of *Cryptosporidium* in primates and herbivores at the Barcelona zoo: a long-term study. Veterinary Parasitology 104: 19–26. PMID: [11779652](#)
33. Onda K, LoBuglio J, Bartram J (2012) Global Access to Safe Water: Accounting for Water Quality and the Resulting Impact on MDG Progress. International Journal of Environmental Research and Public Health 9: 880–894. doi: [10.3390/ijerph9030880](#) PMID: [22690170](#)
34. Akinbo FO, Okaka CE, Omorogie R, Dearen T, Leon ET, et al. (2010) Molecular Characterization of *Cryptosporidium* spp. in HIV-infected Persons in Benin City, Edo State, Nigeria. Fooyin Journal of Health Sciences 2: 85–89.
35. Bridgman SA, Robertson RM, Syed Q, Speed N, Andrews N, et al. (1995) Outbreak of cryptosporidiosis associated with a disinfected groundwater supply. Epidemiology and Infection 115: 555–566. PMID: [8557088](#)
36. Nizeyi JB, Mwebe R, Nanteza A, Cranfield MR, Kalema G, et al. (1999) *Cryptosporidium* sp and *Giardia* sp infections in mountain gorillas (*Gorilla gorilla beringei*) of the Bwindi Impenetrable National Park, Uganda. Journal of Parasitology 85: 1084–1088. PMID: [10647041](#)
37. Graczyk TK, DaSilva AJ, Cranfield MR, Nizeyi JB, Kalema G, et al. (2001) *Cryptosporidium parvum* Genotype 2 infections in free-ranging mountain gorillas (*Gorilla gorilla beringei*) of the Bwindi Impenetrable National Park, Uganda. Parasitology Research 87: 368–370. PMID: [11403378](#)
38. Salzer JS, Rwego IB, Goldberg TL, Kuhlenschmidt MS, Gillespie TR (2007) *Giardia* sp and *Cryptosporidium* sp infections in primates in fragmented and undisturbed forest in western Uganda. Journal of Parasitology 93: 439–440. PMID: [17539436](#)
39. Johnston AR, Gillespie TR, Rwego IB, McLachlan TLT, Kent AD, et al. (2010) Molecular Epidemiology of Cross-Species *Giardia duodenalis* Transmission in Western Uganda. Plos Neglected Tropical Diseases 4(5): e683. doi: [10.1371/journal.pntd.0000683](#) PMID: [20485494](#)
40. Rwego IB, Gillespie TR, Isabirye-Basuta G, Goldberg TL (2008) High rates of *Escherichia coli* transmission between livestock and humans in rural Uganda. Journal of Clinical Microbiology 46: 3187–3191. doi: [10.1128/JCM.00285-08](#) PMID: [18685012](#)
41. Rwego IB, Isabirye-Basuta G, Gillespie TR, Goldberg TL (2008) Gastrointestinal Bacterial Transmission among Humans, Mountain Gorillas, and Livestock in Bwindi Impenetrable National Park, Uganda. Conservation Biology 22: 1600–1607. doi: [10.1111/j.1523-1739.2008.01018.x](#) PMID: [18717695](#)
42. Goldberg TL, Gillespie TR, Rwego IB, Estoff EL, Chapman CA (2008) In Forest Fragmentation as Cause of Bacterial Transmission among Primates, Humans, and Livestock, Uganda. Emerging Infectious Diseases 14(9): 1375–1382. doi: [10.3201/eid1409.071196](#) PMID: [18760003](#)
43. Baldursson S, Karanis P (2011) Waterborne transmission of protozoan parasites: Review of worldwide outbreaks—An update 2004–2010. Water Research 45: 6603–6614. doi: [10.1016/j.watres.2011.10.013](#) PMID: [22048017](#)
44. Aksoy U AC, Sahin S, Usluca S, Yalcin G, Kuralay F, Oral A (2007) First reported waterborne outbreak of cryptosporidiosis with Cyclospora co-infection in Turkey. Euro Surveillance 12 (7):pii = 3142. PMID: [17370959](#)
45. Persson K SB, de Jong B (2007) An outbreak of cryptosporidiosis at a day-care centre in Sweden. European Surveillance 12 (34):pii = 3255.
46. Graczyk TK, Cranfield MR, Fayer R (1997) Recovery of waterborne oocysts of *Cryptosporidium* from water samples by the membrane-filter dissolution method. Parasitology Research 83: 121–125. PMID: [9039693](#)
47. Jiang J, Alderisio KA, Xiao L (2005) Distribution of *Cryptosporidium* Genotypes in Storm Event Water Samples from Three Watersheds in New York. Applied and Environmental Microbiology 71: 4446–4454. PMID: [16085835](#)
48. Hansen JS, Ongerth JE (1991) Effects of time and watershed characteristics on the concentration of *Cryptosporidium* oocysts in river water. Applied and Environmental Microbiology 57: 2790–2795. PMID: [1746940](#)
49. Davies CM, Ferguson CM, Kaucner C, Krogh M, Altavilla N, et al. (2004) Dispersion and transport of *Cryptosporidium* oocysts from fecal pats under simulated rainfall events. Applied and environmental microbiology 70: 1151–1159. PMID: [14766600](#)
50. Xiao LH, Feng YY (2008) Zoonotic cryptosporidiosis. Fems Immunology and Medical Microbiology 52: 309–323. doi: [10.1111/j.1574-695X.2008.00377.x](#) PMID: [18205803](#)

51. Adamu H, Petros B, Zhang G, Kassa H, Amer S, et al. (2014) Distribution and Clinical Manifestations of *Cryptosporidium* Species and Subtypes in HIV/AIDS Patients in Ethiopia. Plos Neglected Tropical Diseases 8(4): e2831. doi: [10.1371/journal.pntd.0002831](https://doi.org/10.1371/journal.pntd.0002831) PMID: [24743521](#)
52. Rajendran P, Ajjampur SSR, Chidambaram D, Kattula D, Rajan DP, et al. (2011) Short Report: Investigation of Potential Zoonotic Transmission of Cryptosporidiosis in Southern India. American Journal of Tropical Medicine and Hygiene 85: 657–659. doi: [10.4269/ajtmh.2011.10-0673](https://doi.org/10.4269/ajtmh.2011.10-0673) PMID: [21976568](#)
53. Abeywardena H, Jex AR, Nolan MJ, Haydon SR, Stevens MA, et al. (2012) Genetic characterisation of *Cryptosporidium* and *Giardia* from dairy calves: Discovery of species/genotypes consistent with those found in humans. Infection Genetics and Evolution 12: 1984–1993. doi: [10.1016/j.meegid.2012.08.004](https://doi.org/10.1016/j.meegid.2012.08.004) PMID: [22981927](#)
54. Xiao LH, Bern C, Arrowood M, Sulaiman I, Zhou L, et al. (2002) Identification of the *Cryptosporidium* pig genotype in a human patient. Journal of Infectious Diseases 185: 1846–1848. PMID: [12085341](#)
55. Wang L, Zhang H, Zhao X, Zhang L, Zhang G, et al. (2013) Zoonotic *Cryptosporidium* Species and *Enterocytozoon bieneusi* Genotypes in HIV-Positive Patients on Antiretroviral Therapy. Journal of Clinical Microbiology 51: 557–563. doi: [10.1128/JCM.02758-12](https://doi.org/10.1128/JCM.02758-12) PMID: [23224097](#)
56. Leoni F, Amar C, Nichols G, Pedraza-Diaz S, McLauchlin J (2006) Genetic analysis of *Cryptosporidium* from 2414 humans with diarrhoea in England between 1985 and 2000. Journal of Medical Microbiology 55: 703–707. PMID: [16687587](#)
57. Li W, Kiulia NM, Mwenda JM, Nyachieo A, Taylor MB, et al. (2011) *Cyclospora papionis*, *Cryptosporidium hominis*, and Human-Pathogenic *Enterocytozoon bieneusi* in Captive Baboons in Kenya. Journal of Clinical Microbiology 49: 4326–4329. doi: [10.1128/JCM.05051-11](https://doi.org/10.1128/JCM.05051-11) PMID: [21956988](#)
58. Leav BA, Mackay MR, Anyanwu A, O'Connor RM, Cevallos AM, et al. (2002) Analysis of sequence diversity at the highly polymorphic Cpgp40/15 locus among *Cryptosporidium* isolates from human immunodeficiency virus-infected children in South Africa. Infection and Immunity 70: 3881–3890. PMID: [12065532](#)